

FINAL EXAM (sample)

NAME(use CAPITAL letters, *first name first*):_____

NAME(sign):_____

SECTION:_____

ID#:_____

Instructions: Each of the 6 problems has worth either 33 or 34 points. Read each question carefully and answer it in the space provided. **YOU MUST SHOW ALL YOUR WORK TO RECEIVE FULL CREDIT.** Clarity of your solutions may be a factor when determining credit. Calculators, books or notes are not allowed. You can use the Formula Sheet provided at our course website. Unless directed to do so, do *not* evaluate complicated expressions to give the result as a fraction or a decimal number. The proctor has been directed not to answer any interpretation questions.

Make sure that you have a total of 8 pages (including this one) with 6 problems.

1	
2	
3	
4	
5	
6	
TOTAL	over 200

Before starting: Imagine that this is your exam and solve it in 2 hours. Problems 1, 2 and 3 come with solutions, so solve them on a different paper. **Avoid reading the solutions**, and compare them with yours only when you finish your simulation.


1. The pair (X, Y) of random variables has ^{joint} density given by ~~$f(x, y)$~~ $f(x, y) = \begin{cases} c(xy + 1) & \text{if } x, y \in [0, 1] \\ 0 & \text{otherwise} \end{cases}$

(a) Determine c . (Recall $\int_0^1 x^n dx = 1/(n+1)$ for $n > -1$.)

$$c \int_0^1 dx \int_0^1 (xy + 1) dy = c \int_0^1 dx (x \cdot \frac{1}{2} + 1)$$

$$= c \left(\frac{1}{2} \cdot \frac{1}{2} + 1 \right) = c \cdot \frac{5}{4} \quad \underline{\underline{c = \frac{4}{5}}}$$

(b) Determine $P(2Y \leq X | Y \leq X)$. = $\frac{P(Y \leq \frac{X}{2})}{P(Y \leq X)}$



$$= \frac{\frac{4}{5} \int_0^1 dx \int_0^{x/2} (xy + 1) dy}{\frac{4}{5} \int_0^1 dx \int_0^x (xy + 1) dy} = \frac{\int_0^1 dx (x \cdot \frac{x^2}{8} + \frac{x}{2})}{\int_0^1 dx (x \cdot \frac{x^2}{2} + x)}$$

$$= \frac{\frac{1}{8} \cdot \frac{1}{4} + \frac{1}{2} \cdot \frac{1}{2}}{\frac{1}{2} \cdot \frac{1}{4} + \frac{1}{2}} = \frac{\frac{1}{16} + \frac{1}{2}}{\frac{1}{4} + 1} = \frac{9}{20}$$

(c) Determine the density of the random variable $Z = \sqrt{X}$.

$$P(Z \leq z) = P(X \leq z^2) = \frac{4}{5} \int_0^{z^2} dx \int_0^1 (xy + 1) dy$$

$$z \in [0, 1]$$

$$= \frac{4}{5} \int_0^{z^2} \left(\frac{1}{2}x + 1 \right) dx$$

$$f_Z(z) = \frac{4}{5} \left(\frac{1}{2} z^2 + 1 \right) \cdot 2z = \frac{4}{5} (z^3 + 2z) \quad \text{if } z \in [0, 1]$$

$$= 0 \quad \text{otherwise}$$

2. Shuffle a full deck of 52 cards.

(a) What is the probability that all Aces are together in the deck (i.e., the four Aces are four consecutive cards, in any order)?

$$\frac{4! \cdot 49!}{52!} \quad \left(\text{or} \quad \frac{49}{\binom{52}{4}} \right)$$

(b) What is the probability that all the Aces are together in the deck and so are Kings, Queens, and Jacks.

$$\frac{(4!)^4 \cdot 40!}{52!}$$

(c) What is the probability that all the Aces are together in the deck and so are all the hearts (♥) cards?



$$52 - 16 = 36 \text{ other cards}$$

$$\frac{2 \cdot 3! \cdot 12! \cdot 37!}{52!}$$

Problem 2, continued.

(d) What is the probability that the first four cards are all hearts?

$$\frac{1}{4} \cdot \frac{12}{51} \cdot \frac{11}{50} \quad \left(\text{or} \quad \frac{\binom{13}{3}}{\binom{52}{3}} \right)$$

(e) What is the probability that the third card is of different suit than either of the first two cards?
(For example, this event happens if the suits are, in order, $\heartsuit\heartsuit\diamondsuit$ or $\heartsuit\spadesuit\diamondsuit$, but not in the case $\heartsuit\spadesuit\heartsuit$.)

$P(\text{1st two cards same suit, 3rd different suit})$

+ $P(\text{1st three cards diff. suits})$

$$= \underset{\substack{\uparrow \\ \text{choose} \\ \text{suit} \\ \text{of 1st 2 cards}}}{4} \cdot \frac{1}{4} \cdot \frac{12}{51} \cdot \frac{39}{50} + \underbrace{4 \cdot 3 \cdot 2}_{\substack{\uparrow \\ \text{suits of} \\ \text{1st 3 cards}}} \cdot \frac{1}{4} \cdot \frac{13}{51} \cdot \frac{13}{50}$$

3. Roll five red dice and five blue dice. All dice are fair.

(a) Let X be the number of 6's rolled on red dice and Y the total number of 6's. Identify the probability mass functions of X and Y . Compute $\text{Var}(Y)$.

X is Binomial $(5, \frac{1}{6})$

Y is Binomial $(10, \frac{1}{6})$

$$\text{Var}(Y) = 10 \cdot \frac{1}{6} \cdot \frac{5}{6} = \frac{50}{6}$$

(b) Let X and Y be as in (a). Determine the joint probability mass function of X and Y . (Give a formula rather than a table.)

$$P(X=x, Y=y) = \underbrace{\binom{5}{x}}_{\text{positions of 6's}} \underbrace{\binom{5}{y-x}}_{\text{other nos.}} 5^{10-y} \cdot \frac{1}{6^{10}}$$

$x = 0, \dots, 5$
 $y = 0, \dots, 5$
 $x \leq y \leq x+5$

(c) Give the conditional probability $P(X=1|Y=1)$ as a single fraction. Compute also the unconditional probability $P(X=1)$. Are X and Y independent?

$$P(X=1|Y=1) = \frac{1}{2} \quad (\text{the 6 needs to appear on a red die})$$

$$P(X=1) = 5 \cdot \frac{1}{6} \cdot \left(\frac{5}{6}\right)^4 = \frac{5^5}{6^5}$$

No. The two are different.

4. Suppose that you toss 2 fair coins. Let X be the number of heads on the first coin and Y the total number of heads.

(a) Compute the joint density of X, Y . Are X and Y independent?

(b) Find $\rho(X, Y)$ and $\mathbb{E}(X|Y)$.

(c) Calculate the characteristic function $\phi_W(t)$, where $W = Y + 1$.

5. Assume that X_1 and X_2 are independent $Exp(1)$. Let $Y_1 = X_1 + X_2$ and $Y_2 = X_1/X_2$.

(a) Compute the joint density of Y_1, Y_2 .

(b) Find F_{Y_2} and $\mathbb{P}(Y_2 \geq 1)$.

(c) Find f_{Y_1} and $\mathbb{E}(Y_1|Y_2)$.

6. Determine whether the following statements are True or False (proof or counterexample).

(a) If $X \sim \text{Ber}(p)$ is independent of $Y \sim \text{Bin}(n, p)$, then $X + Y \sim \text{Bin}(n + 1, p)$.

(b) If $\mathbb{E}(X) = 2$ and $\mathbb{E}(X^2) = 4$, then $\mathbb{E}[(X - 2)^2] = 2$.

(c) If X and Y are independent $\text{Ber}(1/2)$, then $X|(X + Y = 1) \sim \text{Ber}(1/2)$.

(d) If $X \sim \beta(2, 2)$, then $\mathbb{E}(X^n) = \frac{1}{(n + 1)!}$ for all $n \in \mathbb{N}$.

(e) If $f_X(x) = \frac{1}{2}e^{-|x|}$, for $x \in \mathbb{R}$, then the characteristic function of X is $\phi_X(t) = \frac{1}{t^2 + i}$.