

1.4.9. (c) If $ab > 0$ and $bc < 0 \Rightarrow ax^2 + bx + c = 0$ has 2 real sols.

Sketch: $a, b, c \in \mathbb{Z}$, $x \in \mathbb{R}$. We know that the roots are

$$x_{\pm} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{If } b^2 - 4ac = 0 \Rightarrow x_+ = x_- \text{ we do not want it.}$$

If $b^2 - 4ac < 0 \Rightarrow x_+, x_- \in \mathbb{C} - \mathbb{R}$. we do not want it neither.

• If $b^2 - 4ac > 0 \Rightarrow x_+, x_- \in \mathbb{R}$ and $x_+ \neq x_-$. (2 sols).

P: $ab > 0$ and $bc < 0 \Rightarrow \dots \Rightarrow ? \quad ? \Rightarrow \dots \Rightarrow Q$

$$\Rightarrow (ab)(bc) < 0 \Rightarrow b^2 ac < 0 \quad (\Delta)$$

Is it possible $b = 0$? No, if $b = 0 \Rightarrow b^2 ac = 0$.

$$\text{So } b^2 > 0 \text{ by } (\Delta) \quad \frac{b^2 ac}{b^2} < \frac{0}{b^2} \Leftrightarrow ac < 0 \Leftrightarrow 4ac < 0 \Leftrightarrow -4ac > 0$$

Proof: let $a, b, c \in \mathbb{Z}$ and $x \in \mathbb{R}$ such that

$$ab > 0 \text{ and } bc < 0 \Rightarrow (ab)(bc) < 0 \Rightarrow b^2 ac < 0$$

Note that $b = 0 \Rightarrow b^2 ac = 0$. Therefore $b \neq 0$, so

$$b^2 > 0 \Rightarrow \frac{b^2 ac}{b^2} < \frac{0}{b^2} \text{ that means } ac < 0 \Rightarrow -4ac > 0.$$

Now, $b^2 > 0$ and $-4ac > 0$ implies $\overset{\uparrow}{\text{Since } -4 < 0}$

$$b^2 + (-4ac) > 0, \text{ thus } b^2 - 4ac > 0.$$

This means that the discriminant of the eq $ax^2 + bx + c = 0$ is positive so the solutions

$$x_{\pm} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \text{ are real and } x_+ \neq x_-$$

Therefore, the equation has 2 real roots. #

1.5.3 (f) $\exists x, y$ is odd \Rightarrow both x and y are odd
 $P(x, y)$ $Q(x, y)$

Have to do is to show $\sim Q \Rightarrow \sim P$.

Q : " x is odd and y is odd"

$\sim Q$: " x is even or y is even" $(R \vee S) \Rightarrow \sim P$

Case 1: $R \Rightarrow \sim P$

Case 2: $S \Rightarrow \sim P$.

... **DO IT!**

1.6.7i $(\forall \varepsilon > 0)(\exists K \in \mathbb{N})(\forall x \in \mathbb{R})(x > K \Rightarrow \frac{1}{4x} < \varepsilon)$

Proof: Fix $\varepsilon > 0$. Then we can take $K \in \mathbb{N}$

$$K > \frac{1}{4\varepsilon} \Rightarrow \frac{1}{4K} < \varepsilon.$$

Now fix $x \in \mathbb{R}$ s.t. $x > K \Rightarrow$... finish it!

#

want $\frac{1}{4K} < \varepsilon \Leftrightarrow \frac{1}{4\varepsilon} < K$ \leftarrow
Sketch $x > K \Rightarrow 4x > 4K \Rightarrow \frac{1}{4x} < \frac{1}{4K}$

1.6.5(a) $x \in \mathbb{N}$. x prime $\Leftrightarrow \left[x > 1 \wedge \sim \left(\begin{array}{l} \exists k \in \{2, 3, \dots, \} \\ k \leq \sqrt{x} \wedge k | x \end{array} \right) \right]$

Proof: (\Rightarrow) Easy.

$$(\forall k \in \{2, 3, \dots, \}) (k > \sqrt{x} \vee k \nmid x)$$

(\Leftarrow) Fix $x \in \mathbb{N}$.
 x prime iff $(\forall d \in \mathbb{N}) (d | x \Rightarrow (d = 1 \vee d = x))$

Assume that $\forall k \in \{2, 3, \dots, \}$ either $\underbrace{k > \sqrt{x}}_R$ or $\underbrace{k \nmid x}_S$.
 let us prove that x is prime.

Do THE FULL PROOF BY using sketch. //

$$\sim (P \Rightarrow Q) \Leftrightarrow P \wedge \sim Q$$

Sketch: (By contr), $\exists l \in \mathbb{N} \ l | x \wedge \sim (l = 1 \vee l = x)$
 $l | x \wedge l \neq 1 \wedge l \neq x \Leftrightarrow \underbrace{l | x}_S \wedge l \geq 2 \wedge \underbrace{l \neq x}_S$
 $\exists m \in \mathbb{N}$ s.t. $x = l \cdot m \geq l$ so $l \in \{2, 3, \dots, x-1\}$

$\underbrace{l > \sqrt{x}}_0 \Rightarrow \frac{x}{l} = m \neq 1$ since $x \neq l$ $\frac{x}{l} \in \{2, 3, \dots, \}$

either $\underbrace{\frac{x}{l} > \sqrt{x}}_0$ or $\underbrace{\frac{x}{l} \nmid x}_F$ $\frac{x}{l} | x$

so $\underbrace{\frac{x}{l} > \sqrt{x}}_0$ multipl. $l \cdot \frac{x}{l} > \sqrt{x} \cdot \sqrt{x}$


so $x > x$. $(\rightarrow \leftarrow)$

2.3.1k. Find the union and intersection of $\mathcal{A} := \{A_n; n \geq 3\}$
 $A_n = \left[\frac{1}{n}, 2 + \frac{1}{n}\right]$ for $n \in \mathbb{N} - \{1, 2\}$.

Obs: $n=1$ $A_1 = [1, 3]$, $n=2$ $A_2 = \left[\frac{1}{2}, \frac{5}{2}\right]$, $n=3$ $A_3 = \left[\frac{1}{3}, \frac{7}{3}\right]$

$n=4$, $A_4 = \left[\frac{1}{4}, 2 + \frac{1}{4}\right]$

$n=5$ $A_5 = \left[\frac{1}{5}, 2 + \frac{1}{5}\right]$



Guess: (i) $\bigcup_{n \geq 3} A_n = (0, 7/3]$ and (ii) $\bigcap_{n \geq 3} A_n = \left[\frac{1}{3}, 2\right]$

Proof (i) " \subseteq " Pick $x \in \bigcup_{n \geq 3} A_n$, so $\exists n \in \mathbb{N}, n \geq 3$ s.t. $x \in A_n$
 so $x \in \left[\frac{1}{n}, 2 + \frac{1}{n}\right]$ for some $n \geq 3$, and $\left[\frac{1}{n}, 2 + \frac{1}{n}\right] \subseteq (0, 7/3]$
 $\forall n \geq 3$, since $0 < \frac{1}{n}$ and $2 + \frac{1}{n} \leq \frac{7}{3} \quad \forall n \geq 3$.
 Thus $x \in (0, 7/3]$.

" \supseteq " Pick $x \in \mathbb{R}$, $x \in (0, 7/3]$, so either $x \in [1, 7/3]$ or $x \in (0, 1)$. Case 1: $x \in [1, 7/3] \subseteq A_3$ so $x \in A_3 \subseteq \bigcup_{n \geq 3} A_n$ ✓

Case 2: $x \in (0, 1)$, in part $x > 0$ so by AP VII below
 $\exists n \in \mathbb{N}$ s.t. $\frac{1}{n} \leq x$. So if $n \geq 3 \Rightarrow x \in A_3 \subseteq \bigcup_{n \geq 3} A_n$ ✓

And if $n=1$ or $2 \Rightarrow x \geq 1$ or $x \geq 2$ ($\rightarrow \leftarrow$)

In any case $x \in \bigcup_{n \geq 3} A_n$. #

Archimedean Property Version II (AP VII): For every $\varepsilon > 0$
 there exists $n \in \mathbb{N}$ such that $\frac{1}{n} \leq \varepsilon$.

By contradiction, assume $\sim (\forall \varepsilon > 0)(\exists n \in \mathbb{N})\left(\frac{1}{n} \leq \varepsilon\right)$ so
 $(\exists \hat{\varepsilon} > 0)(\forall n \in \mathbb{N})\left(\frac{1}{n} > \hat{\varepsilon}\right)$ so $n < \frac{1}{\hat{\varepsilon}}$, then $\mathbb{N} \rightarrow \leftarrow$
 $\hat{\varepsilon} \in \mathbb{R}$

Because \mathbb{N} is not bounded from above.

Note that if $\varepsilon \in \mathbb{N}$ then AP VII coincides with our
 Arch. Prop. seen in class for $a=1, b=\varepsilon, s=n$.

(i) let's show that $\bigcap_{n \geq 3} A_n = [\frac{1}{3}, 2]$. $A_n = [\frac{1}{n}, 2 + \frac{1}{n}]$

" \supseteq " Since $\frac{1}{n} \leq \frac{1}{3} \forall n \geq 3$, and $2 \leq 2 + \frac{1}{n}, \forall n \geq 3$.

$\Rightarrow \underbrace{[\frac{1}{3}, 2]}_B \subseteq A_n, \forall n \geq 3$ so by Theorem 2.3.2(a)
we conclude that $B = [\frac{1}{3}, 2] \subseteq \bigcap_{n \geq 3} A_n$.

" \subseteq " Pick $x \in \bigcap_{n \geq 3} A_n$, so $x \in A_n = [\frac{1}{n}, 2 + \frac{1}{n}], \forall n \geq 3$

so $\frac{1}{n} \leq x \leq 2 + \frac{1}{n}, \forall n \geq 3$. In particular ($n=3$)

$\frac{1}{3} \leq x$. Also $x \leq 2 + \frac{1}{n}, \forall n \in \mathbb{N}$ (2 options):

1st Say taking $n \rightarrow \infty$ we get $x \leq 2$.
calculus sequences $a_n \leq b_n, \forall n \Rightarrow \lim a_n \leq \lim b_n$ if \lim exists.


2nd: By contradiction, if $x > 2$, $\varepsilon = x - 2 > 0$, --
use AP VII to get $(\rightarrow \leftarrow)$ Thus $x \leq 2$.

therefore $\frac{1}{3} \leq x \leq 2$, so $x \in [\frac{1}{3}, 2]$. $\#$

4.5.10f. HINT: $f: A \rightarrow B, D \subseteq A$

w.t.s. $D = f^{-1}(f(D)) \Leftrightarrow f(A-D) \subseteq B - f(D)$

1st: $D \subseteq f^{-1}(f(D))$ always.
this is item (e).



(\Leftarrow) w.t.s. $f^{-1}(f(D)) \subseteq D$

$x \in f^{-1}(f(D)) \Leftrightarrow f(x) \in f(D) \Rightarrow \boxed{x \in D}$ w.t.s.
In general $x \in D \Rightarrow f(x) \in f(D)$ BUT NOT in general unless f is 1-1.

We know that $f^{-1}(f(D)) \subseteq A$ so $x \in A$.

Assume by contr. $x \notin D \Rightarrow x \in A \wedge x \notin D \Rightarrow$
 $x \in A - D \Rightarrow \dots$ FINISH IT!

4.5.13 $f: A \rightarrow B, x \in A \Rightarrow f(A-x) = f(A) - f(x)$

Proof: RHS \subseteq LHS "always"

Pick $y \in B$ s.t. $y \in f(A) - f(x)$

$\Rightarrow y \in f(A)$ and $y \in f(x)^c$


$\Rightarrow y = f(a)$ for some $a \in A$.

\langle w.t.s. $b \in f(A-x) \Leftrightarrow a \notin X$, so that $a \in A-x \rangle$

By contrad. Assume $a \in X \Rightarrow f(a) \in f(x) \Rightarrow y \in f(x) \rightarrow \in$

so $a \notin X \Rightarrow a \in A-x$ and $y = f(a)$, that


means $y \in f(A-x)$. \langle we did not use 1-1 here \rangle #



S.1. 18(a) HINT: A, B finite $A \approx B$. $f: A \xrightarrow{1-1} B \Rightarrow f$ onto B .


Sketch: If f is not onto B , $\exists b \in B$ s.t. $b \notin \text{Rng}(f)$.

$\exists l \in \mathcal{N}, B \approx \mathcal{N}_l$



On the other hand $A \approx B \Rightarrow \exists g: A \xrightarrow{1-1} B$

By Corollary 4.4.3 $f^{-1}: \text{Rng}(f) \xrightarrow{1-1} A$ is a function



$g \circ f^{-1}: \text{Rng}(f) \xrightarrow{1-1} B \xrightarrow{\varphi} \mathcal{N}_l$

$\exists h: \text{Rng}(f) \xrightarrow{1-1} \mathcal{N}_l$ and show that

for $C = \text{Rng}(f)$, since $b \notin C \xrightarrow{\text{identify } \mathcal{L} \text{ with } \mathcal{L} \setminus \{b\}}$

$\exists \tilde{f}: C \xrightarrow{1-1} \mathcal{N}_{l-1}$ so $\exists \tilde{h}: \mathcal{N}_l \xrightarrow{1-1} \mathcal{N}_{l-1} (\rightarrow \leftarrow)$ by P.P.
 $\tilde{f} \circ h^{-1}$