

1.4.9. (c) If $ab > 0$ and $bc < 0 \Rightarrow ax^2 + bx + c = 0$ has 2 real sols.

Sketch: $a, b, c \in \mathbb{Z}$, $x \in \mathbb{R}$. We know that the roots are

$$x_{\pm} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad \text{If } b^2 - 4ac = 0 \Rightarrow x_+ = x_- \text{ we do not want it.}$$

If $b^2 - 4ac < 0 \Rightarrow x_+, x_- \in \mathbb{C} - \mathbb{R}$. we do not want it neither.

• If $b^2 - 4ac > 0 \Rightarrow x_+, x_- \in \mathbb{R}$ and $x_+ \neq x_-$. (2 sols).

P: $ab > 0$ and $bc < 0 \Rightarrow \dots \Rightarrow ? \quad ? \Rightarrow \dots \Rightarrow Q$

$$\Rightarrow (ab)(bc) < 0 \Rightarrow b^2 ac < 0 \quad (\Delta)$$

Is it possible $b = 0$? No, if $b = 0 \Rightarrow b^2 ac = 0$.

$$\text{So } b^2 > 0 \text{ by } (\Delta) \quad \frac{b^2 ac}{b^2} < \frac{0}{b^2} \Leftrightarrow ac < 0 \Leftrightarrow 4ac < 0 \Leftrightarrow -4ac > 0$$

Proof: let $a, b, c \in \mathbb{Z}$ and $x \in \mathbb{R}$ such that

$$ab > 0 \text{ and } bc < 0 \Rightarrow (ab)(bc) < 0 \Rightarrow b^2 ac < 0$$

Note that $b = 0 \Rightarrow b^2 ac = 0$. Therefore $b \neq 0$, so

$$b^2 > 0 \Rightarrow \frac{b^2 ac}{b^2} < \frac{0}{b^2} \text{ that means } ac < 0 \Rightarrow -4ac > 0.$$

Now, $b^2 > 0$ and $-4ac > 0$ implies $\overset{\text{Since } -4 < 0}{-4ac > 0}$

$$b^2 + (-4ac) > 0, \text{ thus } b^2 - 4ac > 0.$$

This means that the discriminant of the eq $ax^2 + bx + c = 0$ is positive so the solutions

$$x_{\pm} = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \text{ are real and } x_+ \neq x_-$$

Therefore, the equation has 2 real roots. #

1.5.3 (f) $\exists x, y$ is odd \Rightarrow both x and y are odd
 $P(x, y)$ $Q(x, y)$

Have to do is to show $\sim Q \Rightarrow \sim P$.

Q : " x is odd and y is odd"

$\sim Q$: " x is even or y is even" $(R \vee S) \Rightarrow \sim P$

Case 1: $R \Rightarrow \sim P$

Case 2: $S \Rightarrow \sim P$.

... **DO IT!**

1.6.7i $(\forall \varepsilon > 0)(\exists K \in \mathbb{N})(\forall x \in \mathbb{R})(x > K \Rightarrow \frac{1}{4x} < \varepsilon)$

Proof: Fix $\varepsilon > 0$. Then we can take $K \in \mathbb{N}$

$$K > \frac{1}{4\varepsilon} \Rightarrow \frac{1}{4K} < \varepsilon.$$

Now fix $x \in \mathbb{R}$ s.t. $x > K \Rightarrow$... finish it!

#

want $\frac{1}{4K} < \varepsilon \Leftrightarrow \frac{1}{4\varepsilon} < K$ \leftarrow
Sketch $x > K \Rightarrow 4x > 4K \Rightarrow \frac{1}{4x} < \frac{1}{4K}$

1.6.5(a) $x \in \mathbb{N}$. x prime $\Leftrightarrow \left[x > 1 \wedge \sim \left(\begin{array}{l} \exists k \in \{2, 3, \dots, \} \\ k \leq \sqrt{x} \wedge k | x \end{array} \right) \right]$

Proof: (\Rightarrow) Easy.
 (\Leftarrow) Fix $x \in \mathbb{N}$.
 x prime iff $(\forall d \in \mathbb{N}) (d | x \Rightarrow (d = 1 \vee d = x))$
 $(\forall k \in \{2, 3, \dots, \}) (k > \sqrt{x} \vee k \nmid x)$

Assume that $\forall k \in \{2, 3, \dots, \}$ either $k > \sqrt{x}$ or $k \nmid x$.
 let us prove that x is prime. R S

Do THE FULL PROOF BY using sketch. //

Sketch: (By contr), $\exists l \in \mathbb{N} \ l | x \wedge \sim (l = 1 \vee l = x)$
 $l | x \wedge l \neq 1 \wedge l \neq x \Leftrightarrow l | x \wedge l \geq 2 \wedge l \neq x$
 $\exists m \in \mathbb{N}$ s.t. $x = l \cdot m \geq l$ so $l \in \{2, 3, \dots, x-1\}$

$l > \sqrt{x}$ $\frac{x}{l} = m \neq 1$ since $x \neq l$ $\frac{x}{l} \in \{2, 3, \dots, \}$
 either $\frac{x}{l} > \sqrt{x}$ or $\frac{x}{l} \nmid x$ F $\frac{x}{l} | x$


so $\frac{x}{l} > \sqrt{x}$ multipl. $l \cdot \frac{x}{l} > \sqrt{x} \cdot \sqrt{x}$
 so $x > x$. $(\rightarrow \leftarrow)$

2.3.1k. Find the union and intersection of $\mathcal{A} := \{A_n; n \geq 3\}$
 $A_n = \left[\frac{1}{n}, 2 + \frac{1}{n}\right]$ for $n \in \mathbb{N} - \{1, 2\}$.

Obs: $n=1$ $A_1 = [1, 3]$, $n=2$ $A_2 = \left[\frac{1}{2}, \frac{5}{2}\right]$, $n=3$ $A_3 = \left[\frac{1}{3}, \frac{7}{3}\right]$

$n=4$, $A_4 = \left[\frac{1}{4}, 2 + \frac{1}{4}\right]$

$n=5$ $A_5 = \left[\frac{1}{5}, 2 + \frac{1}{5}\right]$



Guess: (i) $\bigcup_{n \geq 3} A_n = (0, 7/3]$ and (ii) $\bigcap_{n \geq 3} A_n = \left[\frac{1}{3}, 2\right]$

Proof (i) " \subseteq " Pick $x \in \bigcup_{n \geq 3} A_n$, so $\exists n \in \mathbb{N}, n \geq 3$ s.t. $x \in A_n$
 so $x \in \left[\frac{1}{n}, 2 + \frac{1}{n}\right]$ for some $n \geq 3$, and $\left[\frac{1}{n}, 2 + \frac{1}{n}\right] \subseteq (0, 7/3]$
 $\forall n \geq 3$, since $0 < \frac{1}{n}$ and $2 + \frac{1}{n} \leq \frac{7}{3} \quad \forall n \geq 3$.
 Thus $x \in (0, 7/3]$.

" \supseteq " Pick $x \in \mathbb{R}$, $x \in (0, 7/3]$, so either $x \in [1, 7/3]$ or $x \in (0, 1)$. Case 1: $x \in [1, 7/3] \subseteq A_3$ so $x \in A_3 \subseteq \bigcup_{n \geq 3} A_n$ ✓

Case 2: $x \in (0, 1)$, in part $x > 0$ so by AP VII below
 $\exists n \in \mathbb{N}$ s.t. $\frac{1}{n} \leq x$. So if $n \geq 3 \Rightarrow x \in A_3 \subseteq \bigcup_{n \geq 3} A_n$ ✓
 And if $n=1$ or $2 \Rightarrow x \geq 1$ or $x \geq 2$ ($\rightarrow \leftarrow$)

In any case $x \in \bigcup_{n \geq 3} A_n$. #

Archimedean Property Version II (AP VII): For every $\varepsilon > 0$
 there exists $n \in \mathbb{N}$ such that $\frac{1}{n} \leq \varepsilon$.

By contradiction, assume $\sim (\forall \varepsilon > 0)(\exists n \in \mathbb{N})\left(\frac{1}{n} \leq \varepsilon\right)$ so
 $(\exists \hat{\varepsilon} > 0)(\forall n \in \mathbb{N})\left(\frac{1}{n} > \hat{\varepsilon}\right)$ so $n < \frac{1}{\hat{\varepsilon}}$, then $\mathbb{N} \rightarrow \leftarrow$
 $\hat{\varepsilon} \in \mathbb{R}$

Because \mathbb{N} is not bounded from above.

Note that if $\varepsilon \in \mathbb{N}$ then AP VII coincides with our
 Arch. Prop. seen in class for $a=1, b=\varepsilon, s=n$.

(i) let's show that $\bigcap_{n \geq 3} A_n = [\frac{1}{3}, 2]$. $A_n = [\frac{1}{n}, 2 + \frac{1}{n}]$

" \supseteq " Since $\frac{1}{n} \leq \frac{1}{3} \forall n \geq 3$, and $2 \leq 2 + \frac{1}{n}, \forall n \geq 3$.

$\Rightarrow \underbrace{[\frac{1}{3}, 2]}_B \subseteq A_n, \forall n \geq 3$ so by Theorem 2.3.2(a)
we conclude that $B = [\frac{1}{3}, 2] \subseteq \bigcap_{n \geq 3} A_n$.

" \subseteq " Pick $x \in \bigcap_{n \geq 3} A_n$, so $x \in A_n = [\frac{1}{n}, 2 + \frac{1}{n}], \forall n \geq 3$

so $\frac{1}{n} \leq x \leq 2 + \frac{1}{n}, \forall n \geq 3$. In particular ($n=3$)

$\frac{1}{3} \leq x$. Also $x \leq 2 + \frac{1}{n}, \forall n \in \mathbb{N}$ (2 options):

1st Say taking $n \rightarrow \infty$ we get $x \leq 2$.
calculus sequences $a_n \leq b_n, \forall n \Rightarrow \lim a_n \leq \lim b_n$ if \lim exists.

2nd: By contradiction, if $x > 2$, $\varepsilon = x - 2 > 0$, --
use AP VII to get $(\rightarrow \leftarrow)$ Thus $x \leq 2$.

therefore $\frac{1}{3} \leq x \leq 2$, so $x \in [\frac{1}{3}, 2]$. $\#$